



DISKRETE MATHEMATIK UND OPTIMIERUNG

Luis A. Goddyn
Winfried Hochstättler:

Nowhere-zero flows in regular matroids and Hadwiger's Conjecture

Technical Report feu-dmo031.13
Contact: goddyn@sfu.ca
winfried.hochstaettler@fernuni-hagen.de

FernUniversität in Hagen
Fakultät für Mathematik und Informatik
Lehrgebiet für Diskrete Mathematik und Optimierung
D – 58084 Hagen

2000 Mathematics Subject Classification: 05C15, 05B35, 52C40

Keywords: nowhere zero flow, regular matroid, chromatic number, flow number, total unimodularity

NOWHERE-ZERO FLOWS IN REGULAR MATROIDS AND HADWIGER'S CONJECTURE

LUIS A. GODDYN AND WINFRIED HOCHSTÄTTLER

ABSTRACT. We present a tool that shows, that the existence of a k -nowhere-zero-flow is compatible with 1-,2- and 3-sums in regular matroids. As application we present a conjecture for regular matroids that is equivalent to Hadwiger's conjecture for graphs and Tutte's 4- and 5-flow conjectures.

KEYWORDS: *nowhere zero flow, regular matroid, chromatic number, flow number, total unimodularity*

1. INTRODUCTION

A (real) matrix is *totally unimodular* (TUM) if each subdeterminant belongs to $\{0, \pm 1\}$. Totally unimodular matrices enjoy several nice properties which give them a fundamental role in combinatorial optimization and matroid theory. In this note we prove that the TUM possesses an attractive property.

Let $S \subseteq \mathbb{R}$, and let A be a real matrix. A column vector f is a S -flow of A if $Af = 0$ and every entry of f is a member of $\pm S$.

For any additive abelian group Γ use the notation $\Gamma^* = \Gamma \setminus \{0\}$. For a TUM A and a column vector f with entries in Γ , the product Af is a well defined column vector with entries in Γ , by interpreting $(-1)\gamma$ to be the additive inverse of γ .

It is convenient to use the language of matroids. A regular oriented matroid M is an oriented matroid that is representable $M = M[A]$ by a TUM matrix A . Here the elements $E(M)$ of M label the columns of A . Each (signed) cocircuit $D = (D^+, D^-)$ of M corresponds to a $\{0, \pm 1\}$ -valued vector in the row space of A and having minimal support. The $+1$ -entries in this vector constitute the sets D^+ . It is known [19, Prop. 1.2.5] that two TUMs represent the same oriented matroid if and only if the first TUM can be converted to the second TUM by a succession of the following operations: multiplying a row by -1 , adding one row to another, deleting a row of zeros, and permuting columns (with their labels).

For $S \subseteq E(M)$ we use the notation $f(S) = \sum_{e \in S} f(e)$. Let $M = M[A]$ be the regular oriented matroid represented by the TUM A . Let $S \subseteq \Gamma$ where Γ is an abelian group. An S -flow of M is a function $f : E(M) \rightarrow S$ for which $Af = 0$, where f is interpreted to be a vector indexed by the column labels of A . For any $S \subseteq \Gamma$ we say that a regular matroid M has an S -flow if any of the TUMs that represent M has an S -flow. By the previous paragraph, this property of M is well defined. Since the rows of a TUM A generate the cocycle space of $M = M[A]$, we have that a function $f : E(M) \rightarrow \Gamma$ is a flow if and only if for every signed cocircuit $D = (D^+, D^-)$ we have that $f(D) = 0$ where $f(D)$ is defined to equal $f(D^+) - f(D^-)$.

Let Γ be a finite abelian group. Let M be a regular oriented matroid, and let $F \subseteq E(M)$ and let $f : F \rightarrow \Gamma$. Let $\tau_\Gamma(M, f)$ denote the number of Γ^* -flows of M which are extensions of f .

THEOREM 1. *Let M be an regular oriented matroid. Let $F \subseteq E(M)$ and let $f, f' : F \rightarrow \Gamma$. Suppose that for every minor N of M satisfying $E(N) = F$, we have that f is a Γ -flow of N if and only if f' is a Γ -flow of N . Then $\tau_\Gamma(M, f) = \tau_\Gamma(M, f')$.*

Proof. We proceed by induction on $d = |E \setminus F|$. If $d = 0$, then there is nothing to prove. Otherwise let $e \in E \setminus F$. If e is a coloop of M , then $\tau_\Gamma(M, f) = \tau_\Gamma(M, f') = 0$. If e is a loop of M , then by applying induction to $M \setminus e$, we have $\tau_\Gamma(M, f) = \tau_\Gamma(M, f') = (|\Gamma| - 1)\tau_\Gamma(M \setminus e, f)$. Otherwise we apply Tutte's deletion/contraction formula [3] and induction to get

$$\tau_\Gamma(M, f') = \tau_\Gamma(M/e, f') - \tau_\Gamma(M \setminus e, f') = \tau_\Gamma(M/e, f) - \tau_\Gamma(M \setminus e, f) = \tau_\Gamma(M, f).$$

□

COROLLARY 2. *Let D be a positively oriented cocircuit of a regular oriented matroid M . Let $f, f' : D \rightarrow \Gamma$. Suppose that for every $S \subseteq D$ we have that $f(S) = 0$ if and only if $f'(S) = 0$. Then $\tau_\Gamma(M, f) = \tau_\Gamma(M, f')$.*

Proof. Let N be a minor of M satisfying $E(N) = D$. Then $E(N)$ is a disjoint union $\bigcup_i D_i$ of positively oriented cocircuits of N [9, Prop. 9.3.1]. Thus f is a Γ^* -flow of N if and only if f has no zeros, and $f(D_i) = 0$ for each i . The result follows from Theorem 1. \square

COROLLARY 3. *Let M be a regular oriented matroid which has a Γ^* -flow f .*

- (1) *Let $e \in E(M)$ and $\gamma \in \Gamma^*$. Then M has a Γ^* -flow f' with $f'(e) = \gamma$.*
- (2) *Let D be a signed cocircuit of M of cardinality three. Let $f' : D \rightarrow \Gamma^*$ satisfy $f'(D) = 0$. Then f' extends to a Γ^* -flow of M .*

Proof. (1) In any minor N with $E(N) = \{e\}$, both f' and $f \upharpoonright_{\{e\}}$ are Γ^* -flows of N if and only if N is a loop. Thus by Theorem 1 $\tau_\Gamma(M, f') = \tau_\Gamma(M, f) > 0$.
(2) Let $S \subset D$. For any $e \in D$ we have $f'(D \setminus \{e\}) = f'(D) - f'(e) = -f'(e) \neq 0$. Therefore $f'(S) = 0$ if and only if $S = D$. Since f is a Γ -flow and D is a positively oriented cocircuit of D we have $f(D) = 0$. Since $f(e) \neq 0$ for $e \in D$ we again have that $f(S) = 0$ if and only if $S = D$. It follows from Theorem 1 that $\tau_\Gamma(M, f') = \tau_\Gamma(M, f) > 0$. \square

A k -nowhere zero flow (k -NZF) of a regular oriented matroid M is an S -flow of M for $S = \{1, 2, \dots, k-1\} \subset \mathbb{R}$. We frequently use the following observation of Tutte [15].

PROPOSITION 4. *Let Γ be an abelian group of order k , and let $S = \{1, 2, \dots, k-1\} \subset \mathbb{R}$. Then M has a k -NZF if and only if M has a Γ^* -flow. In particular, the existence of a Γ^* -flow in M depends only on $|\Gamma|$.*

A key step in the proof of Proposition 4 is the conversion of a Γ^* -flow into a k -NZF, where Γ is the group of integers modulo k . By modifying this argument, one can show that the statement of Corollary 3 remains true if each occurrence of the symbol Γ^* is replaced by the set of integers $S = \{\pm 1, \pm 2, \dots, \pm(k-1)\}$. We omit the proof of this fact, as it is not needed in this paper.

2. SEYMOUR DECOMPOSITION

We provide here a description of Seymour's decomposition theorem for regular oriented matroids. We refer the reader to [13] for further details. We first describe three basic types of regular oriented matroids.

A oriented matroid is *graphic* if it can be represented by the $\{0, \pm 1\}$ -valued vertex-edge incidence matrix of a directed graph, where loops and multiple edges are allowed. Any $\{0, \pm 1\}$ -valued matrix whose rows span the nullspace of a network matrix is called a *dual network matrix*. Dual network matrices are also TUM, and an oriented matroid is *cographic* if it is representable by a dual network matrix. The third class consists of all the all the orientations of one special regular matroid R_{10} . Every orientation of R_{10} can be represented by the matrix $[I|B]$ where B is obtained by negating a subset of the columns of the following matrix.

$$(1) \quad \begin{bmatrix} + & 0 & 0 & + & - \\ - & + & 0 & 0 & + \\ + & - & + & 0 & 0 \\ 0 & + & - & + & 0 \\ 0 & 0 & + & - & + \end{bmatrix}$$

Here “+” and “−” respectively denote +1 and −1.

Let M_1, M_2 be regular oriented matroids. If $E(M_1)$ and $E(M_2)$ are disjoint, then the 1-sum $M_1 \oplus_1 M_2$ is just the direct sum of M_1 and M_2 . The signed cocircuits of $M_1 \oplus_1 M_2$ are the signed subsets of $E(M_1) \cup E(M_2)$ which are signed cocircuits of either M_1 or M_2 . If $M_1 \cap M_2 = \{e\}$ and e is neither a loop nor a coloop in each M_i , then the 2-sum $M_1 \oplus_2 M_2$ has element set $E(M_1) \Delta E(M_2)$, where “ Δ ” is the symmetric difference operator. A signed cocircuit is a signed subset of $E(M_1 \oplus_2 M_2)$ that is either a signed cocircuit of M_1 or M_2 , or is a signed set of the form

$$(2) \quad D = (D_1^+ \Delta D_2^+, D_1^- \Delta D_2^-)$$

where each (D_i^+, D_i^-) is a signed cocircuit of M_i , and $e \in (D_1^+ \cap D_2^+) \cup (D_1^- \cap D_2^-)$. If $M_1 \cap M_2 = B$ and $B = (B^+, B^-)$ is a signed cocircuit of cardinality 3 in each M_i , then the 3-sum $M_1 \oplus_3 M_2$ has element set $E(M_1) \Delta E(M_2)$. A signed cocircuit is a signed subset of $E(M_1 \oplus_3 M_2)$ that is either a signed cocircuit of M_1 or M_2 , or a signed subset of the form (2) where each (D_i^+, D_i^-) is a signed cocircuit of M_i , with $D_1 \cap D_2 = \emptyset$ and (B^+, B^-) equals one of the following ordered pairs:

$$\begin{aligned} & ((D_1^+ \cap B^+) \cup (D_2^+ \cap B^+), (D_1^- \cap B^-) \cup (D_2^- \cap B^-)) \\ & ((D_1^- \cap B^+) \cup (D_2^- \cap B^+), (D_1^+ \cap B^-) \cup (D_2^+ \cap B^-)). \end{aligned}$$

The oriented version of Seymour's decomposition theorem [13] and can be derived from [5, Theorem 6.6].

THEOREM 5. *Every regular oriented matroid M can be constructed by means of repeated application of k -sums, $k = 1, 2, 3$, starting with oriented matroids, each of which is isomorphic to a minor of M and each of which is either graphic, cographic, or an orientation of R_{10} .*

We note that Schriver [12] states an equivalent version of Theorem 5 in terms of TUMs, that requires a second representation of R_{10} in (1) due to his implicit selection of a basis.

Here is the main tool of this paper, which we employ in two subsequent applications.

THEOREM 6. *Let $k \geq 2$ be an integer and let \mathcal{M} be a set of regular oriented matroids that is closed under minors. If every graphic and cographic member of \mathcal{M} has a k -NZF, then every matroid in \mathcal{M} has a k -NZF.*

Proof. Let $M \in \mathcal{M}$. We proceed by induction on $|E(M)|$. If M is an orientation of R_{10} , then M has a 2-NZF since R_{10} is a disjoint union of circuits, and each circuit is the support of a $\{0, \pm 1\}$ -flow in M . If M is graphic or cographic, then we are done by assumption. Otherwise, by Theorem 5, M has two proper minors $M_1, M_2 \in \mathcal{M}$ such that $M = M_1 \oplus_i M_2$, for some $i = 1, 2, 3$. By induction, each M_i has a k -NZF. Thus by Proposition 4, both minors have a Γ^* -flow where Γ is any fixed group of order k . By Corollary 3, we may assume that these Γ^* -flows coincide on $M_1 \cap M_2$. Hence the union of these functions is a well defined Γ^* -flow on M and we are done by another application of Proposition 4. \square

3. TUTTE'S FLOW CONJECTURES AND HADWIGER'S CONJECTURE

In this section we will present a conjecture that unifies two of Tutte's Flow Conjectures and Hadwiger's Conjecture on graph colorings.

CONJECTURE 7 (H(k)[4]). *If a simple graph is not k -colorable, then it must have a K_{k+1} -minor.*

While H(1) and H(2) are trivial, Hadwiger proved his conjecture for $k = 3$ and pointed out that Klaus Wagner proved that H(4) is equivalent to the Four Color Theorem [18, 2, 10]. Robertson, Seymour and Thomas [11] reduced H(5) to the Four Color Theorem. The conjecture remains open for $k \geq 6$.

Tutte [15] pointed out that the Four Color Theorem is equivalent to the statement that every planar graph admits an 4-NZ-flow. Generalizing this to arbitrary graphs he conjectured that

CONJECTURE 8 (Tutte's Flow Conjecture [15]). *There is a finite number $k \in \mathbb{N}$ such that every bridgeless graph admits a k -NZ-flow.*

and moreover that

CONJECTURE 9 (Tutte's Five Flow Conjecture [15]). *Every bridgeless graph admits a 5-NZ-flow.*

Note that the latter is best possible as the Petersen graph does not admit a 4-NZ-flow. Conjecture 8 has been proven independently by Kilpatrick [7] and Jaeger [6] with $k = 8$ and improved to $k = 6$ by Seymour [14].

Conjecture 9 has a sibling which is a more direct generalization of the Four Color Theorem.

CONJECTURE 10 (Tutte's Four Flow Conjecture [16, 17]). *Every graph without a Petersen-minor admits a 4-NZ-flow.*

In [16, 17] Tutte cited Hadwiger's conjecture as a motivating theme and pointed out that while

“Hadwiger's conjecture asserts that the only irreducible chain-group which is graphic is the coboundary group of the complete 5-graph”

Conjecture 10 means that

“the only irreducible chain-group which is cographic is the cycle group of the Petersen graph.”

The first statement refers to the case where the rows of a totally unimodular matrix A consist of a basis of signed characteristic vectors of cycles of a digraph.

Combining these we derive the following formulation in terms of regular matroids. First let us call any integer combination of the rows of A a *coflow*. Clearly, by duality resp. orthogonality, flows and coflows yield the same concept in regular matroids. Note that the existence of a k -NZ-coflow in a graph is equivalent to k -colorability [16].

CONJECTURE 11 (Tutte’s Four Flow Conjecture, matroid version). *A regular matroid that does not admit a 4-NZ-flow has either a minor isomorphic to the cographic matroid of the K_5 or a minor isomorphic to the graphic matroid of the Petersen graph.*

Equivalently, we have

CONJECTURE 12 (Hadwiger’s Conjecture for regular matroids and $k = 4$). *A regular matroid that is not 4-colorable, i.e. that does not admit a NZ-4-coflow, has a K_5 or a Petersen-dual as a minor.*

Some progress concerning this Conjecture was made by Lai, Li and Poon using the Four Color Theorem

THEOREM 13 ([8]). *A regular matroid that is not 4-colorable has a K_5 or a K_5 -dual as a minor.*

Tutte’s Five Flow Conjecture now suggests the following matroid version of Hadwiger’s conjecture:

CONJECTURE 14 (Hadwiger’s Conjecture for regular matroids and $k \geq 5$). *If a regular matroid is not k -colorable for $k \geq 5$, then it must have a K_{k+1} -minor.*

THEOREM 15. (1) *Conjecture 11 is equivalent to Conjecture 10.*

(2) *Conjecture 14 for $k = 5$ is equivalent to Conjecture 9.*

(3) *Conjecture 14 for $k \geq 6$ is equivalent to Conjecture 7.*

Proof. (1) By Weiske’s Theorem [4] a graphic matroid has no K_5^* -minor. Hence Conjecture 11 clearly implies Conjecture 10. The other implication is proven by induction on $|E(M)|$. Consider a regular matroid M , that is not 4-colorable, i.e. that does not admit a NZ-4-coflow. Clearly, M cannot be isomorphic to R_{10} . If M is graphic, it must have a K_5 -minor by the Four Color Theorem [2, 10] and an observation of Klaus Wagner [18]. If M is cographic it must have a Petersen-dual-minor by Conjecture 10. Otherwise, by Theorem 5, M has two proper minors $M_1, M_2 \in \mathcal{M}$. such that $M = M_1 \oplus_i M_2$, for some $i = 1, 2, 3$ and at least one of them is not 4-colorable by Theorem 6. Using induction we find either a Petersen-dual-minor or a K_5 -minor in one of the M_i and hence also in M . Thus, Conjecture 10 implies Conjecture 11.

(2) We proceed as in the first case using $H(5)$ for graphs [11] instead of the Four Color Theorem.

(3) We proceed similar to the first case, with only a slight difference in the base case. If M is graphic, it must have a K_{k+1} -minor by Conjecture 7. M cannot be cographic by Seymour’s 6-flow-theorem [14]. \square

REMARK 16. *James Oxley pointed that Theorem 15 could also be proven using splitting formulas for the Tutte polynomial (see e.g. [1]), Seymour’s decomposition and the fact that the flow number as well as the chromatic number are determined by the smallest non-negative integer non-zero of certain evaluations of the Tutte polynomial.*

REFERENCES

1. Artur Andrzejak, *Splitting formulas for tutte polynomials*, Journal of Combinatorial Theory, Series B **70** (1997), no. 2, 346–366.
2. Kenneth I. Appel and Wolfgang Haken, *Every planar map is four colorable*, Bull. Amer. Math. Soc. **82** (1976), no. 5, 711–712.
3. D.K Arrowsmith and F Jaeger, *On the enumeration of chains in regular chain-groups*, Journal of Combinatorial Theory, Series B **32** (1982), no. 1, 75–89.
4. Hugo Hadwiger, *Über eine Klassifikation der Streckenkomplexe*, Vierteljahresschrift der Naturforschenden Gesellschaft in Zürich **88** (1943), 133–142.

5. Winfried Hochstättler and Robert Nickel, *The flow lattice of oriented matroids*, Contributions to Discrete Mathematics **2** (2007), no. 1, 68–86.
6. F. Jaeger, *Flows and generalized coloring theorems in graphs*, Journal of Combinatorial Theory, Series B **26** (1979), no. 2, 205 – 216.
7. Peter Allan Kilpatrick, *Tutte's first colour-cycle conjecture.*, Master's thesis, University of Cape Town, 1975.
8. Hong-Jian Lai, Xiangwen Li, and Hoifung Poon, *Nowhere zero 4-flow in regular matroids*, J. Graph Theory **49** (2005), no. 3, 196–204.
9. James G. Oxley, *Matroid theory*, The Clarendon Press Oxford University Press, New York, 1992.
10. Neil Robertson, Daniel Sanders, Paul Seymour, and Robin Thomas, *The Four-Colour theorem*, Journal of Combinatorial Theory, Series B **70** (1997), no. 1, 2–44.
11. Neil Robertson, Paul Seymour, and Robin Thomas, *Hadwiger's conjecture for k_6 -free graphs*, Combinatorica **13** (1993), 279–361.
12. Alexander Schrijver, *Theory of linear and integer programming*, Wiley, June 1998.
13. P. D. Seymour, *Decomposition of regular matroids*, J. Combin. Theory Ser. B **28** (1980), no. 3, 305–359. MR 579077 (82j:05046)
14. ———, *Nowhere-zero 6-flows*, J. Combin. Theory Ser. B **30** (1981), no. 2, 130–135. MR MR615308 (82j:05079)
15. W.T. Tutte, *A contribution to the theory of chromatic polynomials*, Canad. J. Math. **6** (1954), 80–91.
16. ———, *On the algebraic theory of graph colorings*, Journal of Combinatorial Theory **1** (1966), no. 1, 15 – 50.
17. ———, *A geometrical version of the four color problem*, Combinatorial Math. and Its Applications (R. C. Bose and T. A. Dowling, eds.), Chapel Hill, NC: University of North Carolina Press, 1967.
18. K. Wagner, *Über eine Eigenschaft der ebenen Komplexe*, Mathematische Annalen **114** (1937), no. 1, 570–590.
19. Neil White, *Combinatorial geometries*, Cambridge University Press, September 1987.